

ON THE CONNECTION BETWEEN SETS OF OPERATOR SYNTHESIS AND SETS OF SPECTRAL SYNTHESIS FOR LOCALLY COMPACT GROUPS

JEAN LUDWIG AND LYUDMILA TUROWSKA

We extend the results by Froelich and Spronk & Turowska on the connection between operator synthesis and spectral synthesis for $A(G)$ to second countable locally compact groups G . This gives us another proof that one-point subset of G is a set of spectral synthesis and that any closed subgroup is a set of local spectral synthesis. Furthermore we show that “non-triangular” sets are strong operator Ditkin sets and we establish a connection between operator Ditkin sets and Ditkin sets. These results are applied to prove that any closed subgroup of G is a local Ditkin set.

1. Introduction

In [A] Arveson discovered a connection between the invariant subspace theory and spectral synthesis. He defined (operator) synthesis for subspace lattices and proved the failure of operator synthesis by using the famous example of Schwartz on non-synthesizability of the two-sphere S^2 for $A(\mathbb{R}^3)$. In [F] Froelich made this connection more precise for separable abelian group. For G a separable compact group this relation was obtained in [ST, Theorem 4.6.]. We generalize these results to second countable locally compact groups (Theorems 4.3 and 4.10). We use the definition of sets of operator synthesis as defined in [ShT1]. We prove that a closed subset $E \subset G$ is set of local spectral synthesis for $A(G)$ if and only if the diagonal set $E^* = \{(s, t) \in G \times G \mid st^{-1} \in E\}$ is a set of operator synthesis with respect to Haar measure. We remark that if $A(G)$ has an (unbounded) approximate identity then any set of local spectral synthesis is a set of synthesis for $A(G)$ and any compact set of local spectral synthesis is globally spectral for any group G . We give a new proof that a one-point set is spectral and any closed subgroup of second countable group is a set of local spectral synthesis. Using operator synthesizability of sets of finite width we obtain certain example of sets of spectral synthesis.

Operator-Ditkin sets have been defined in [ShT1]. In Section 5 we show first that “non-triangular” type sets are strong operator-Ditkin. The proof is inspired by [D]. For G second countable group we prove that the diagonal set $E^* \subset G \times G$ is a strong operator-Ditkin set with respect to Haar measure then $E \subset G$ is a local Ditkin set for $A(G)$ and conversely, for any strong Ditkin set $E \subset G$, the set E^* is operator-Ditkin with respect to Haar measure. As an application we obtain that any closed subgroup of a second countable group is a local Ditkin set. This result was known for neutral subgroups of arbitrary locally compact groups G ([DD]) and amenable groups G ([FKLS]).

2. Preliminaries and notations

Let G be a locally compact σ -compact separable group with left Haar measure $m = dg$. Let $L^p(G)$, $p = 1, 2$, denote the space of p -integrable functions with norm $\|\cdot\|_p$ and let $C_c(G)$ denote the algebra of continuous compactly supported complex-valued functions on G . The convolution algebra $L^1(G)$ is an involutive algebra with involution defined by $f^*(s) = \Delta^{-1}(s)f(s^{-1})$, where Δ is the modulus of the group. Let Σ be the set of all (equivalence classes of) continuous unitary representations π of G in Hilbert spaces H_π . For $f \in L^1(G)$, $\pi \in \Sigma$, we put $\pi(f) = \int_G f(g)\pi(g)dg \in B(H_\pi)$ with the integral converging in the strong operator topology, and then

$$\|f\|_\Sigma = \sup_{\pi \in \Sigma} \|\pi(f)\|,$$

where $\|\cdot\|$ is the operator norm in $B(H_\pi)$.

The *enveloping C^* -algebra* $C^*(G)$ of G is the completion of $L^1(G)$ with respect to $\|\cdot\|_\Sigma$. Let $\lambda : G \rightarrow B(L_2(G))$ be the left regular representation given by $\lambda(s)f(g) = f(s^{-1}g)$. We denote by $C_r^*(G)$ the *reduced C^* -algebra* of G , that is the C^* -algebra generated by operators $\lambda(f) \in B(L_2(G))$, $f \in L^1(G)$, and by $VN(G)$ the von Neumann algebra of G , that is

$$VN(G) = \overline{\text{span}}^{WOT} \{\lambda(g) : g \in G\} = \overline{C_r^*(G)}^{WOT} \subset B(L_2(G)).$$

The *Fourier-Stieltjes algebra*, $B(G)$, is the set of all coefficients $s \mapsto \pi_{\xi, \eta}(s) = (\pi(s)\xi, \eta)$, where $\pi \in \Sigma$, $\xi, \eta \in H_\pi$, of unitary representations of G , as defined by Eymard, [E]. $B(G)$ is a Banach algebra with respect to the norm

$$\|u\| = \inf\{\|\xi\| \|\eta\| : u = \pi_{\xi, \eta}\}.$$

Note that $B(G) \simeq C^*(G)^*$.

The *Fourier algebra*, $A(G)$, is the family of functions $s \mapsto (\lambda(s)\xi, \eta) = \bar{\eta} * \check{\xi}$, $\xi, \eta \in L_2(G)$, $\check{\xi}(s) = \xi(s^{-1})$. $A(G)$ is identified with the predual $VN(G)_*$ via $\langle (\lambda(s)\xi, \eta), T \rangle = (T\xi, \eta)$, and thus is a normed algebra with the norm denoted by $\|\cdot\|_A$. It is known that for $u \in A(G)$ there exist even $\xi, \eta \in L_2(G)$ such that $u = \bar{\eta} * \check{\xi}$ and $\|u\|_A = \|\xi\|_2 \cdot \|\eta\|_2$. Furthermore, $A(G)$ is a closed ideal in $B(G)$.

Since $A(G)$ is the predual of the von Neumann algebra $VN(G)$, $A(G)$ possesses a structure of operator space and one can define a notion of completely bounded multipliers, $M_{cb}A(G)$, for $A(G)$. For the theory of operator space and completely bounded maps we refer the reader to [Efr, BSm, S]. A complex-valued function $u : G \rightarrow \mathbb{C}$ is a completely bounded multiplier if it is a multiplier, i.e. $uA(G) \subset A(G)$, and is completely bounded as a linear map on $A(G)$. We have $A(G) \subset B(G) \subset M_{cb}A(G)$ and, for $u \in A(G)$, $\|u\|_{cb} \leq \|u\|_A$, where $\|\cdot\|_{cb}$ is the completely bounded norm (see [S, Corollary 2.3.3]). We will use here a characterization of $M_{cb}A(G)$ obtained by N.Spronk in [S] as formulated in Theorem 3.1.

3. Spectral and operator synthesis

Let A be a semisimple, regular, commutative Banach algebra with X_A as spectrum; for any $a \in A$ we shall denote then by $\hat{a} \in C_0(X_A)$ its Gelfand transform. Let also $E \subset X_A$ be a closed subset. We then denote by

$$I_A(E) = \{a \in A \mid \hat{a}^{-1}(0) \text{ contains } E\},$$

$$J_A^0(E) = \{a \in A \mid \hat{a}^{-1}(0) \text{ contains a neighborhood of } E\} \text{ and } J_A(E) = \overline{J_A^0(E)}.$$

It is known that $I_A(E)$ and $J_A(E)$ are the largest and the smallest closed ideals with E as hull, i.e., if I is a closed ideal such that $\{x \in X_A : f(x) = 0 \text{ for all } f \in I\} = E$ then

$$J_A(E) \subset I \subset I_A(E).$$

Let $I_A^c(E)$ denote the set of all compactly supported functions $f \in I_A(E)$. We say that E is a *set of spectral synthesis* (*local spectral synthesis*) for A if $J_A(E) = I_A(E)$ ($I_A^c(E) \subset J_A(E)$).

Let A^* be the dual of A . For $a \in A$ we set $\text{supp}(a) = \overline{\{x \in X_A : \hat{a}(x) \neq 0\}}$ and $\text{null}(a) = \{x \in X_A : \hat{a}(x) = 0\}$. For $\tau \in A^*$ and $a \in A$ define $a\tau$ in A^* by $a\tau(b) = \tau(ab)$ and define the support of τ by

$$\text{supp}(\tau) = \{x \in X_A : a\tau \neq 0 \text{ whenever } a(x) \neq 0\}.$$

Then, for a closed set $E \subset X_A$

$$J_A(E)^\perp = \{\tau \in A^* : \text{supp}(\tau) \subset E\}$$

and E is spectral for A if and only if $\tau(a) = 0$ for any $a \in A$ and $\tau \in A^*$ such that $\text{supp}(\tau) \subset E \subset \text{null}(a)$.

The algebra $A(G)$ is a semi-simple abelian regular Banach algebras with spectrum G . In what follows we write $I_A(E)$ for $I_{A(G)}(E)$, $J_A(E)$ for $J_{A(G)}(E)$, and $\text{supp}_{VN} T$ for $\text{supp}(T)$ if $T \in VN(G) = (A(G))^*$.

Now we recall some definitions and important facts on operator synthesis following [A, ShT1]. To make use of the results from [A, ShT1] we will assume in the rest of the paper that G is a second countable locally compact group or s.c.l.c group for short, and therefore is metrizable by [HRI, 8.3].

A subset $E \subset G \times G$ is called *marginally null* (with respect to $m \times m$) if $E \subset (M \times G) \cup (G \times N)$ and $m(M) = m(N) = 0$. Two subsets E_1, E_2 are marginally equivalent ($E_1 \sim^M E_2$ or simply $E_1 \cong E_2$) if their symmetric difference is marginally null. Furthermore, $E_1 \subset^M E_2$ means that $E_1 \setminus E_2$ is marginally null, a property holds marginally almost everywhere if it holds everywhere apart of a marginally null set, and so on.

Following [ErKSh] we define a pseudo-topology on G . We call a subset E a pseudo-open if it is marginally equivalent to a countable union of measurable rectangles $A \times B$. The complements of pseudo-open sets are pseudo-closed sets.

Set $T(G) = L^2(G) \hat{\otimes} L^2(G)$, where $\hat{\otimes}$ denotes the projective tensor product. Note that in [ShT1] the notation $\Gamma(G, G)$ is used instead of $T(G)$. Every $\Psi \in T(G)$ can be identified with a function $\Psi : G \times G \rightarrow \mathbb{C}$ which admits a representation

$$(3.1) \quad \Psi(x, y) = \sum_{n=1}^{\infty} f_n(x)g_n(y)$$

where $f_n \in L^2(G)$, $g_n \in L^2(G)$ and $\sum_{n=1}^{\infty} \|f_n\|_2 \cdot \|g_n\|_2 < \infty$. Such a representation defines a function marginally almost everywhere (m.a.e.), so two functions in $T(G)$ which coincides m.a.e. are identified. The $L^2(G) \hat{\otimes} L^2(G)$ -norm of Ψ is

$$\|\Psi\|_{T(G)} = \inf \left\{ \sum_{n=1}^{\infty} \|f_n\|_2 \cdot \|g_n\|_2 : \Psi = \sum_{n=1}^{\infty} f_n \otimes g_n \right\}.$$

Note that a simple renormalization shows that each $\Psi \in T(G)$ admits a representation (3.1) such that $\sum_{n=1}^{\infty} \|f_n\|_2^2 \cdot \sum_{n=1}^{\infty} \|g_n\|_2^2 < \infty$ and the norm $\|\Psi\|_{T(G)}$ can be taken as the

square root of the infimum of $\sum_{n=1}^{\infty} \|f_n\|_2^2 \cdot \sum_{n=1}^{\infty} \|g_n\|_2^2$ over all such representations. By [ErKSh] any $\Psi \in T(G)$ is pseudo-continuous. Thus if Ψ vanishes (m.a.e.) on $K \subset G$ it vanishes on a pseudo-closed set. For $\mathcal{F} \subset T(G)$, the null set, null \mathcal{F} , is defined to be the largest pseudo-closed set such that each function $F \in \mathcal{F}$ vanishes on it. For a pseudo-closed set $E \subset G \times G$, let

$$\Phi(E) = \{w \in T(G) : w = 0 \text{ m.a.e. on } E\},$$

$$\Phi_0(E) = \overline{\{w \in T(G) : w = 0 \text{ on a pseudo-neighbourhood of } E\}}.$$

The spaces $\Phi(E)$, $\Phi_0(E)$ are $L^\infty(G) \times L^\infty(G)$ -bimodules: if $f, g \in L^\infty(G)$ and $w \in \Phi(E)$ ($w \in \Phi_0(E)$) then $f(x)g(y)w(x, y) \in \Phi(E)$ ($f(x)g(y)w(x, y) \in \Phi_0(E)$) respectively). Moreover, $\Phi(E)$ and $\Phi_0(E)$ are the largest and the smallest $L^\infty(G) \times L^\infty(G)$ -invariant subspaces of $T(G)$ whose null set is E .

A subset $E \subset G \times G$ is called a *set of (operator) synthesis* or *synthetic* with respect to m if $\Phi(E) = \Phi_0(E)$.

It is known that $B(L_2(G)) \simeq T(G)^*$ (see [A]). The duality is given by

$$\langle T, \Psi \rangle = \sum_{n=1}^{\infty} (Tf_n, \bar{g}_n),$$

for $T \in B(L_2(G))$ and $\Psi = \sum_{n=1}^{\infty} f_n \otimes g_n \in T(G)$.

Let P_U denote the multiplication operators by the characteristic functions of a subset $U \subset G$. We say that $T \in B(L_2(G))$ is *supported* in $E \subset G \times G$ (or E supports T) if $P_V T P_U = 0$ for each pair of Borel sets $U \subset G, V \subset G$ such that $(U \times V) \cap E \cong \emptyset$. Then there exists the smallest (up to a marginally null set) pseudo-closed set, $\text{supp} T$, which supports T . We write $\text{supp}(T) \subset E$ if T is supported by E . In the seminal paper [A] Arveson defined a support in a similar way but using closed sets instead of pseudo-closed. This closed support, $\text{supp}_A T$ can be strictly larger than $\text{supp}(T)$. Then E is a set of operator synthesis if $\langle T, w \rangle = 0$ for any $T \in B(L_2(G))$ and $w \in T(G)$ with $\text{supp}(T) \subset E \subset \text{null } w$ (the inclusions up to a marginally null set).

We consider also the space $V^\infty(G)$ of all (marginal equivalence classes of) functions $\Psi(x, y)$ that can be written in the form (3.1) with $f_n \in L^\infty(G), g_n \in L^\infty(G)$ and

$$\sum_{n=1}^{\infty} |f_n(x)|^2 \leq C, \quad x \in G, \quad \sum_{n=1}^{\infty} |g_n(y)|^2 \leq C, \quad y \in G,$$

and for such Ψ we have

$$\|\Psi\|_{V^\infty} = \inf \left\{ \left\| \sum_{n=1}^{\infty} |f_n|^2 \right\|_\infty^{1/2} \left\| \sum_{n=1}^{\infty} |g_n|^2 \right\|_\infty^{1/2} : \Psi = \sum_{n=1}^{\infty} f_n \otimes g_n \right\}.$$

In tensor notations $V^\infty(G) = L^\infty(G) \hat{\otimes}^{w^*h} L^\infty(G)$, the weak*-Haagerup tensor product ([BSm]). Let

$$V_{inv}^\infty(G) = \{w \in V^\infty(G) : w(sr, tr) = w(s, t) \text{ for all } r \text{ in } G \text{ and m.a.e. } (s, t) \in G \times G\}.$$

In [S] Spronk found a connection between $V_{inv}^\infty(G)$ and the algebra $M_{cb}A(G)$ of completely bounded multipliers of $A(G)$. For a function $u : G \rightarrow \mathbb{C}$ and $t, s \in G$ define

$$(Nu)(t, s) = u(ts^{-1}).$$

Theorem 3.1. [S] *The map $u \mapsto Nu$ is a complete isometry from $M_{cb}A(G)$ onto $V_{inv}^\infty(G)$.*

4. Spectral synthesis and operator synthesis

In this section we will prove our main result establishing a connection between operator synthesis and spectral synthesis for $A(G)$, where G is a second countable locally compact group. The proofs are inspired by the proof of [ST, Theorem 4.6].

The Banach space $B(L_2(G))$ is a left $V^\infty(G)$ -module with the action defined for $w = \sum_{i=1}^\infty \varphi_i \otimes \psi_i \in V^\infty(G)$ and $T \in B(L_2(G))$ by

$$w \cdot T = \sum_{i=1}^\infty M_{\psi_i} T M_{\varphi_i},$$

where the partial sums converge strongly. The operator $T \mapsto \sum_{i=1}^\infty M_{\psi_i} T M_{\varphi_i}$ we will also denote by Δ_w .

For a closed subset $E \subset G$ we set

$$E^* = \{(s, t) \in G \times G \mid st^{-1} \in E\}.$$

Lemma 4.1. *Let $S \in VN(G)$. Then*

$$\text{supp}(S) \subset \{\text{supp}_{VN} S\}^*.$$

Proof. Let U, V be closed subsets of G such that $(U \times V) \cap \{\text{supp}_{VN}(S)\}^* = \emptyset$. Then there exists an open neighborhood, W , of $\{\text{supp}_{VN} S\}^*$ such that $(U \times V) \cap W = \emptyset$. Take f and g in $L_2(G)$ such that $\text{supp}(f) \subset U$, $\text{supp}(g) \subset V$. For $u = \langle \lambda(\cdot)f, g \rangle$, we have $\langle Sf, g \rangle = \langle S, u \rangle$. Moreover, $\text{supp}(u) \subset UV^{-1}$ and $\text{supp}(u) \cap \text{supp}_{VN} S \subset UV^{-1} \cap \text{supp}_{VN} S = \emptyset$. Thus $0 = \langle S, u \rangle = \langle Sf, g \rangle$. As f and g are chosen arbitrarily, $P_V S P_U = 0$. By the regularity of m , the last holds for any Borel sets U, V giving the statement. \square

Remark 4.2. Let H be a closed subgroup of G . Then

$$\begin{aligned} H^* &= \{(s, t) \in G \times G : st^{-1} \in H\} = \{(s, t) \in G \times G : Hs = Ht\} \\ &= \{(s, t) \in G \times G : f(t) = f(s)\}, \end{aligned}$$

where $f : G \rightarrow H \backslash G$ is a continuous mapping defined by $f(t) = Ht$.

Assume $\text{supp}_{VN}(S) \subset H$. By Lemma 4.1, $\text{supp}(S) \subset H^*$. As for any Borel set $\Delta \subset H \backslash G$ and $\alpha = f^{-1}(\Delta)$, we have $(\alpha^c \times \alpha) \cap H^* = \emptyset$, it gives $P_{\alpha^c} S P_\alpha = 0$. Since this is true for any Δ we have also $P_\alpha S P_{\alpha^c} = 0$ implying that $P_\alpha S = S P_\alpha$ and hence S belongs to the commutant \mathcal{B}' of the von Neumann algebra \mathcal{B} generated by multiplication operators by functions $\varphi \in L^\infty(G)$ which are constant on the right cosets.

Theorem 4.3. *Let G be a s.c.l.c. group and $E \subset G$ be a closed subset. If E^* is synthetic with respect to m then E is a set of local synthesis for $A(G)$.*

Proof. Assume that E^* is synthetic with respect to m . Let $u \in I_A^c(E)$ and $S \in VN(G)$, $\text{supp}_{VN}(S) \subset E$. By Theorem 3.1, $Nu \in V^\infty(G)$. Moreover,

$$(4.2) \quad uT = \Delta_{Nu}T \text{ for any } T \in VN(G).$$

In fact, if $Nu(s, t) = \sum_i \varphi_i(t) \psi_i(s)$, then

$$\begin{aligned} \Delta_{Nu} \lambda(s) f(t) &= \sum_i M_{\varphi_i} \lambda(s) M_{\psi_i} f(t) = \\ &= \sum_i \varphi_i(t) \psi_i(s^{-1}t) f(s^{-1}t) = Nu(t, s^{-1}t) f(s^{-1}t) = u(s) \lambda(s) f(t) \end{aligned}$$

for any $f \in L^2(G)$ and $s \in G$. The operator Δ_{Nu} is weakly-continuous. In fact, if $S_k \rightarrow 0$ weakly, $\|S_k\| \leq C$ for some constant C and

$$\begin{aligned} |(\Delta_{Nu}(S_k)f, g)| &\leq \sum_{i=1}^n |(S_k\psi_i f, \bar{\varphi}_i g)| + \sum_{i=n+1}^{\infty} |(S_k\psi_i f, \bar{\varphi}_i g)| \\ &\leq \sum_{i=1}^n |(S_k\psi_i f, \bar{\varphi}_i g)| + \|S_k\| \left(\sum_{i=n+1}^{\infty} \|\psi_i f\|^2 \right)^{1/2} \left(\sum_{i=n+1}^{\infty} \|\varphi_i g\|^2 \right)^{1/2} \\ &= \sum_{i=1}^n |(S_k\psi_i f, \bar{\varphi}_i g)| + C \left(\int_G \sum_{i=n+1}^{\infty} |\psi_i(t)f(t)|^2 dt \right)^{1/2} \left(\int_G \sum_{i=n+1}^{\infty} |\varphi_i(t)g(t)|^2 dt \right)^{1/2}. \end{aligned}$$

For given $\varepsilon > 0$, by Lebesgue's theorem, there exists n such that the second summand is less than ε and then, as $S_k \rightarrow 0$ weakly, there exists K such that the first summand is less than ε for any $k \geq K$. Therefore (4.2) holds for any $T \in VN(G)$.

Clearly, since $u \in I_A(E)$, Nu vanishes on E^* . By Lemma 4.1, we also have that $\text{supp}(S) \subset \{\text{supp}_{VN} S\}^* \subset E^*$. Therefore, for each $w \in T(G)$,

$$\langle \Delta_{Nu} S, w \rangle = \langle S, (Nu)w \rangle = 0$$

so that $uS = \Delta_{Nu} S = 0$.

From the regularity of $A(G)$ it follows that there exists a compactly supported function $v \in A(G)$ such that $v = 1$ on the support of u . Thus

$$\langle S, u \rangle = \langle S, vu \rangle = \langle uS, v \rangle = 0.$$

□

It is easy to see that the condition for E to be a set of local synthesis for $A(G)$ is equivalent to the condition $uS = 0$ for any $u \in A(G)$ and $S \in VN(G)$ such that $\text{supp}_{VN}(S) \subset E \subset \text{null } u$. If G is amenable then we have the implication

$$(4.3) \quad uT = 0 \Rightarrow \langle T, u \rangle = 0,$$

guaranteed by the existence of a bounded approximate identity and any set of local synthesis is a set of synthesis. Certainly the assumption of boundedness of the identity is superfluous, and the statement holds even for G such that $A(G)$ has an unbounded approximate identity. So that we have

Corollary 4.4. *Let G be a s.c.l.c. group such that $A(G)$ has an (unbounded) approximate identity and let E be a closed subset of G . If E^* is a set of operator synthesis with respect to m then E is a set of spectral synthesis.*

It is not known whether an approximate identity exists for any locally compact group G . Unbounded approximate identities which are completely bounded as multipliers of the Fourier algebra $A(G)$ were studied in [CaH, Ca, CoH]. Those exist for a number of groups like the general Lorentz group $SO_0(n, 1)$, its closed subgroups, in particular, the free group \mathbb{F}_n on n generators, extensions of $SO_0(n, 1)$ by a finite group, in particular, $SL(2, \mathbb{R})$, $SL(2, \mathbb{C})$, $SL(2, \mathbb{H})$, and weakly-amenable groups.

The property (4.3) of an operator $T \in VN(G)$ was discussed in [E] and called there by the property (H). In particular, it was shown that any T supported in a compact set

E possesses this property. Thus any compact set of local synthesis is a set of spectral synthesis.

Corollary 4.5. *Let E be a compact subset of G . If E^* is a set of synthesis with respect to m then E is a set of spectral synthesis.*

It is not known whether there exists $T \in VN(G)$ which does not satisfy (4.3).

The next statement was proved by Eymard for arbitrary locally compact groups G , [E], using more complicated arguments of the theory of distributions on G .

Corollary 4.6. *Let s be an element of the s.c.l.c. group G . Then $\{s\}$ is a set of spectral synthesis.*

Proof. It is enough to prove the statement for $E = \{e\}$, where e is the identity element in G . In this case $E^* = \{(s, s) : s \in G\}$. That E^* is a set of operator synthesis follows e.g. from [ShT1, Theorem 4.8], but it can be easily seen also using the following simple arguments.

Let $T \in B(L_2(G))$ and $\text{supp}(T) \subset E^*$. It follows from Remark 4.2 that T is the multiplication operator by some function $a \in L^\infty(G, m)$. Let now $F = \sum_{i=1}^{\infty} f_i \otimes g_i \in \Phi(E^*)$. Then

$$\langle T, F \rangle = \sum_{i=1}^{\infty} \langle Tg_i, \bar{f}_i \rangle = \sum_{i=1}^{\infty} \langle ag_i, \bar{f}_i \rangle = \int_G a(r)F(r, r)dr = 0,$$

and therefore E^* is a set of synthesis with respect to m . By Corollary 4.5, E is a set of spectral synthesis. \square

We also have another proof in the case of s.c.l.c. groups of the following statement proved by Herz, [He, Theorem 2].

Corollary 4.7. *Any closed subgroup H of G is a set of local spectral synthesis.*

Proof. We have $H^* = \{(s, t) \in G \times G : st^{-1} \in H\} = \{(s, t) \in G \times G : Hs = Ht\} = \{(s, t) \in G \times G : f(t) = f(s)\}$, where $f : G \rightarrow H \setminus G$ is a continuous function defined by $f(t) = Ht$. By [ShT1, Theorem 4.8], H^* is a set of operator synthesis and the statement now follows from Theorem 4.3. \square

Remark 4.8. By Corollaries 4.4, 4.5, any compact subgroup H is spectral for $A(G)$ and any closed subgroup is spectral for $A(G)$ if $A(G)$ has an approximate identity.

That any closed subgroup of a locally compact group G is a set of spectral synthesis was shown by Takesaki and Tatsuuma, [TT, Theorem 3], using the Mackey imprimitivity theorem and the result mentioned in Remark 4.2 proved by a different method.

Example 4.9. Let R be an ordered s.c.l.c. group and $\Delta : G \rightarrow R$ be a continuous homomorphism. Take a finite intersection of intervals $\cap_{k=1}^m [\alpha_k, \beta_k]$ in R and set $E = \Delta^{-1}(\cap_{k=1}^m [\alpha_k, \beta_k])$. Then

$$\begin{aligned} E^* &= \{(s, t) : \alpha_k \leq \Delta(st^{-1}) \leq \beta_k, k = 1, \dots, m\} \\ &= \{(s, t) : \alpha_k \Delta(t) \leq \Delta(s) \leq \beta_k \Delta(t), k = 1, \dots, m\} \\ &= \{(s, t) : f_i^k(t) \leq g_i^k(s), i = 1, 2, k = 1, \dots, m\}, \end{aligned}$$

where $f_1^k(t) = \alpha_k \Delta(t)$, $f_2^k(t) = \Delta(t)^{-1}$, $g_1^k(s) = \Delta(s)$, $g_2^k(s) = \Delta(s)^{-1} \beta_k$. By [ShT1, Theorem 4.8], E^* is a set of operator synthesis with respect to the Haar measure on G . Therefore E is a set of spectral synthesis by Theorem 4.3.

Our next aim is to prove a converse to the statement of Theorem 4.3. The Banach space $T(G)$ is an $L^1(G)$ -module with the action defined by

$$f \odot w(s, t) = \int_G f(r) \Delta^{1/2}(r) w(sr, tr) dr, \quad f \in L^1(G), w \in T(G),$$

where Δ is the modular function of G . We have $\|f \odot w\|_{T(G)} \leq \|f\|_1 \cdot \|w\|_{T(G)}$. Moreover, $e_\alpha \odot w \rightarrow w$ for any bounded approximate identity $\{e_\alpha\}$ in $L^1(G)$ (see [HR11, 32.22, 32.33] and [ST, p.365]).

Let us define another action by compactly supported $L^1(G)$ -functions f :

$$f \cdot w(s, t) = \int_G f(r) w(sr, tr) dr, \quad w \in T(G).$$

Then $f \odot w = f \Delta^{1/2} \cdot w$.

We observe that by the estimate (4.4) below, the integral $\int_G f(r) w(sr, tr) dr, w \in T(G), s, t \in G$, converges also if $f \in L^\infty(G)$ and so defines a mapping $(s, t) \rightarrow f \cdot w(s, t) := \int_G f(r) w(sr, tr) dr$.

Theorem 4.10. *Let G be a s.c.l.c. group. If a closed subset $E \subset G$ is a set of local spectral synthesis for $A(G)$ then E^* is synthetic with respect to Haar measure.*

Proof. It is sufficient to show that $w \cdot T = 0$ for $T \in B(L_2(G))$ and $w \in V^\infty(G)$ such that $\text{supp}(T) \subset E^* \subset \text{null } w$ ([ShT2, Proposition 5.3]).

As G is second countable the group G is σ -compact and therefore there exist compact sets K_n such that $K_n \subset K_{n+1}$ and $\bigcup_{n=1}^\infty K_n = G$. Then clearly, $M_{\chi_{K_n}} T M_{\chi_{K_n}} \rightarrow T$ strongly. Therefore we can restrict ourselves to a compactly supported operator T , i.e. $\text{supp}(T) \subset M \times M$ for a compact set $M \subset G$, and compactly supported $w \in V^\infty(G)$. Note that in this case $w \in T(G)$.

Let \hat{G} denote the set of (equivalence classes of) irreducible continuous unitary representations of G . For $\pi \in \hat{G}$ let H_π denote the representation space of π . Fix a basis $\{e_j\}_j$ in H_π and denote by u_{kj}^π the matrix coefficients of π , i.e. $u_{kj}^\pi(s) = (\pi(s)e_k, e_j)$.

For (a compactly supported) $w = \sum_{i=1}^\infty \varphi_i \otimes \psi_i \in T(G)$ and $\pi \in \hat{G}$ consider now the following operator-valued function

$$w^\pi(s, t) = \int_G w(sr, tr) \pi(r) dr.$$

The integral is well-defined as a Bochner integral. In fact, for each $s, t \in G$, applying Cauchy-Schwartz's inequality, we obtain

$$\begin{aligned}
(4.4) \quad \int_G |w(sr, tr)| dr &\leq \int_G \sum_{i=1}^{\infty} |\varphi_i(sr)\psi_i(tr)| dr \\
&\leq \int_G \left(\sum_{i=1}^{\infty} |\varphi_i(sr)|^2 \right)^{1/2} \left(\sum_{i=1}^{\infty} |\psi_i(tr)|^2 \right)^{1/2} dr \\
&\leq \left(\sum_{i=1}^{\infty} \int_G |\varphi_i(sr)|^2 dr \right)^{1/2} \left(\sum_{i=1}^{\infty} \int_G |\psi_i(tr)|^2 dr \right)^{1/2} \\
&= \left(\sum_{i=1}^{\infty} \|\varphi_i\|_2^2 \right)^{1/2} \left(\sum_{i=1}^{\infty} \|\psi_i\|_2^2 \right)^{1/2} < \infty.
\end{aligned}$$

Set $\tilde{w}^\pi(s, t) = \pi(s)w^\pi(s, t)$ and

$$(4.5) \quad w_{kj}^\pi(s, t) = (w^\pi(s, t)e_k, e_j) = u_{kj}^\pi \cdot w(s, t), \quad \tilde{w}_{kj}^\pi(s, t) = (\tilde{w}^\pi(s, t)e_k, e_j).$$

If $w \in \Phi(E^*)$ then $w(sr, tr)$ vanishes m.a.e. on E^* for all r , and therefore $w^\pi(s, t)$, $\tilde{w}^\pi(s, t)$, $w_{kj}^\pi(s, t)$ and $\tilde{w}_{kj}^\pi(s, t)$ vanish on E^* .

We have the following expression for w_{kj}^π and \tilde{w}_{kj}^π :

$$\begin{aligned}
\tilde{w}_{kj}^\pi(s, t) &= \int_G w(sr, tr)(\pi(sr)e_k, e_j) dr = \int_G w(r, ts^{-1}r)(\pi(r)e_k, e_j) dr \\
&= \int_G w(r, ts^{-1}r)u_{kj}^\pi(r) dr.
\end{aligned}$$

In particular, if $w = f \otimes g \in T(G)$, then

$$(4.6) \quad \tilde{w}_{kj}^\pi(s, t) = \int_G f(r)u_{kj}^\pi(r)g(ts^{-1}r) dr = N(fu_{kj}^\pi * \check{g})(s, t).$$

Furthermore

$$\begin{aligned}
w_{kj}^\pi(s, t) &= \int_G w(sr, tr)(\pi(r)e_k, e_j) dr = \int_G w(r, ts^{-1}r)(\pi(s^{-1}r)e_k, e_j) dr \\
&= \int_G w(r, ts^{-1}r)(\pi(r)e_k, \pi(s)e_j) dr \\
&= \sum_l \int_G w(r, ts^{-1}r)(\pi(r)e_k, e_l)(e_l, \pi(s)e_j) dr \\
(4.7) \quad &= \sum_l u_{lj}^\pi(s) \int_G w(r, ts^{-1}r)u_{kl}^\pi(r) dr = \sum_l u_{lj}^\pi(s)\tilde{w}_{kl}^\pi(s, t).
\end{aligned}$$

We state first that $\tilde{w}_{kj}^\pi(s, t) \in V^\infty(G)$. Indeed, if $w = f \otimes g \in T(G)$, then by (4.6) $\tilde{w}_{kj}^\pi = N(fu_{kj}^\pi * \check{g})$ and therefore, since $fu_{kj}^\pi * \check{g} \in A(G)$, $\tilde{w}_{kj}^\pi \in V^\infty(G)$, by Theorem 3.1. Moreover,

$$\|\tilde{w}_{kj}^\pi(s, t)\|_{V^\infty} \leq \|fu_{kj}^\pi\|_2 \|g\|_2 \leq \|f\|_2 \|g\|_2,$$

so that the linear operator $w \mapsto \tilde{w}_{kj}^\pi \in V^\infty(G)$ defined on elementary tensors extends to a bounded operator $T(G) \rightarrow V^\infty(G)$. Thus $\tilde{w}_{kj}^\pi \in V^\infty(G)$ for any $w \in T(G)$.

Next we show that $w_{kj}^\pi \in V^\infty(G)$. For $T \in B(L^2(G))$ and $w = f \otimes g \in T(G)$ such that $\|w\| = \|f\|_2 \|g\|_2$, define $w_{kj}^\pi \cdot T$ by

$$(4.8) \quad \langle w_{kj}^\pi \cdot T, \Psi \rangle = \sum_l \langle \tilde{w}_{kl}^\pi \cdot T, u_{lj}^\pi(s) \Psi(s, t) \rangle, \quad \Psi \in T(G).$$

This formula makes sense. In fact, if $\Psi = \sum_{i=1}^\infty f_i \otimes g_i$, such that $\|\Psi\|_{T(G)}^2 = \sum_{i=1}^\infty \|f_i\|_2^2 \sum_{i=1}^\infty \|g_i\|_2^2$, we have

$$\begin{aligned} \left| \sum_l \langle \tilde{w}_{kl}^\pi \cdot T, u_{lj}^\pi \Psi \rangle \right|^2 &\leq \left(\sum_l \|\tilde{w}_{kl}^\pi \cdot T\|^2 \right) \left(\sum_l \|u_{lj}^\pi \Psi\|_{T(G)}^2 \right) \\ &\leq \left(\sum_l \|\tilde{w}_{kl}^\pi\|_{V^\infty}^2 \|T\|^2 \right) \left(\sum_l \sum_{i=1}^\infty \|u_{lj}^\pi f_i\|_2^2 \cdot \sum_{i=1}^\infty \|g_i\|_2^2 \right) \\ &\leq \|T\|^2 \left(\sum_l \|f u_{kl}^\pi\|_2^2 \|g\|_2^2 \right) \left(\sum_{i=1}^\infty \sum_l \|u_{lj}^\pi f_i\|_2^2 \right) \cdot \sum_{i=1}^\infty \|g_i\|_2^2 \\ &= \|T\|^2 \|f\|_2^2 \|g\|_2^2 \sum_{i=1}^\infty \|f_i\|_2^2 \cdot \sum_{i=1}^\infty \|g_i\|_2^2 \\ &\leq \|T\|^2 \|w\|_{T(G)}^2 \|\Psi\|_{T(G)}^2, \end{aligned}$$

the last equality follows from

$$\begin{aligned} \sum_l \|f u_{kl}^\pi\|_2^2 &= \sum_l \int_G |f(t) u_{kl}^\pi(t)|^2 dt = \int_G |f(t)|^2 \sum_l |(\pi(t) e_k, e_l)|^2 dt \\ &= \int_G |f(t)|^2 \|(\pi(t) e_k)\|_2^2 dt = \int_G |f(t)|^2 dt = \|f\|_2^2. \end{aligned}$$

Thus the operator $w_{kj}^\pi \cdot T$ is well-defined for $w = f \otimes g$ and since $\|w_{kj}^\pi \cdot T\| \leq \|T\| \|w\|_{T(G)}$ for elementary tensors w , the definition $w_{kj}^\pi \cdot T$ makes sense for any $w \in T(G)$ and

$$(4.9) \quad \left| \sum_l \langle \tilde{w}_{kl}^\pi \cdot T, u_{lj}^\pi \Psi \rangle \right|^2 \leq \|T\|^2 \|w\|_{T(G)}^2 \|\Psi\|_{T(G)}^2$$

for any $\Psi \in T(G)$. Clearly, $T \mapsto w_{kj}^\pi \cdot T$ is thus a bounded $L^\infty(G)$ -bimodule map on $B(L^2(G))$ and hence by [Sm, 2.1] is completely bounded. Then by [BSm, 4.2], it is of the form $\omega \cdot T$ for $\omega \in V^\infty(G)$ and therefore $w_{kj}^\pi = \omega$ m.a.e.

For \tilde{w}^π we have

$$\begin{aligned} \tilde{w}^\pi(sr, tr) &= \pi(sr) w^\pi(sr, tr) = \pi(sr) \int_G w(sr p, tr p) \pi(p) dp \\ &= \pi(s) \int_G w(sr p, tr p) \pi(rp) dp = \pi(s) \int_G w(sp, tp) \pi(p) dp = \tilde{w}^\pi(s, t), \end{aligned}$$

implying $\tilde{w}_{kj}^\pi \in V_{inv}^\infty(G)$. By Theorem 3.1,

$$(4.10) \quad \tilde{w}_{kj}^\pi = Nu$$

for some $u \in M_{cb}A(G)$. Moreover, if w vanishes on E^* then u vanishes on E .

We claim that $Nu \cdot T = 0$ for any operator T and $u \in M_{cb}A(G)$ such that $\text{supp}(T) \subset E^*$, null $u \supset E$. In fact, since E is a set of local spectral synthesis, given $w \in A(G)$ with compact support, uw can be approximated by $u_\alpha \in J_A^0(E)$ and therefore $N(uw)$ is a $V^\infty(G)$ -limit of Nu_α , vanishing on pseudo-neighborhoods of E^* . By [ShT1, Theorem 4.3], $\langle T, \Psi \rangle = 0$ for any $\Psi \in \Phi_0(E^*)$. Therefore $\langle Nu_\alpha \cdot T, F \rangle = \langle T, (Nu_\alpha)F \rangle = 0$ for any $F \in T(G)$ implying $Nu_\alpha \cdot T = 0$ and $N(uw) \cdot T = 0$. As $\langle Nu \cdot T, (Nw)F \rangle = 0$ for any $w \in A(G)$ with compact support and $F \in T(G)$ it is enough to see now that the subspace, \mathcal{M} , generated by $(Nw)F$, where $F \in T(G)$ and $w \in A(G)$ with compact support, is dense in $T(G)$. As null $\mathcal{M} = \emptyset$, this is true by an analogue of Wiener's Tauberian Theorem [ShT1, Corollary 4.3].

We obtain by (4.10) that for any $w \in V^\infty(G)$ which is compactly supported and vanishes on E^*

$$\tilde{w}_{kj}^\pi \cdot T = 0.$$

Therefore by (4.9)

$$w_{kj}^\pi \cdot T = \left(\sum_l u_{jl}^\pi(s) \tilde{w}_{kj}^\pi(s, t) \right) \cdot T = 0,$$

Let K be a compact set such that $\text{supp}(w) \subset K \times K$ and $\text{supp}(T) \subset K \times K$. Then since $T = M_{\chi_K} T M_{\chi_K}$, for compactly supported $h \in L^1(G)$ we have

$$\begin{aligned} \langle T, h \cdot w \rangle &= \langle M_{\chi_K} T M_{\chi_K}, h \cdot w \rangle = \langle T, \chi_K(t) \chi_K(s) \int_G w(sr, tr) h(r) dr \rangle \\ &= \langle T, \int_G w(sr, tr) \chi_{K^{-1}K}(r) h(r) dr \rangle = \langle T, h \chi_{K^{-1}K} \cdot w \rangle. \end{aligned}$$

Take $f \in L^\infty(G)$, $g \in L^\infty(G)$, $\text{supp}(f) \subset K$, $\text{supp}(g) \subset K$, and set $\omega = f \otimes g$. Then

$$\begin{aligned} \langle \omega \cdot T, u_{kj}^\pi \chi_{K^{-1}K} \cdot w \rangle &= \langle T, (u_{kj}^\pi \chi_{K^{-1}K} \cdot w) \omega \rangle \\ &= \langle T, w_{kj}^\pi \omega \rangle = \langle w_{kj}^\pi \cdot T, \omega \rangle = 0. \end{aligned}$$

Hence for finite linear combinations $\sum_i c_i u_i$ of matrix coefficients

$$\langle \omega \cdot T, \sum_i c_i u_i \chi_{K^{-1}K} \cdot w \rangle = 0.$$

Take now an approximate identity $\{e_\alpha\}$ consisting of non-negative continuous functions with compact support in $L^1(G)$. We can assume that $K^{-1}K$ contains the supports of e_α 's. Then, by [Di, 13.6.5], $\Delta^{1/2} e_\alpha$ can be approximated in $L^1(G)$ by finite linear combinations $\sum_{i=1}^d c_i u_i \chi_{K^{-1}K}$. This yields

$$0 = \langle \omega \cdot T, (\Delta^{1/2} e_\alpha) \cdot w \rangle = \langle \omega \cdot T, (e_\alpha) \odot w \rangle$$

and therefore $0 = \langle \omega \cdot T, w \rangle = \langle w \cdot T, \omega \rangle$. Finally, we obtain $w \cdot T = 0$. □

Corollary 4.11. *Let G be a s.c.l.c. group.*

a) *Then a compact set $E \subset G$ is a set of spectral synthesis for $A(G)$ if and only if E^* is a set of operator synthesis with respect to Haar measure.*

b) *Assume that $A(G)$ has an approximate identity. Then a closed set $E \subset G$ is a set of spectral synthesis for $A(G)$ if and only if E^* is a set of operator synthesis with respect to Haar measure.*

Proof. Follows from Corollary 4.4, Corollary 4.5 and Theorem 4.10. □

5. Ditkin sets and operator Ditkin sets.

Our goal in this section is to find a connection between Ditkin sets and operator Ditkin sets. Let G be a locally compact group and let m be the Haar measure on G . A closed subset $E \subset G$ is said to be a (local) Ditkin set if for any $f \in I_A(E)$ ($f \in I_A^c(E)$) there exists a sequence $\{\tau_n\}_{n=1}^\infty$ such that $\tau_n \in J_A^0(E)$ ($n = 1, 2, \dots$) and $\tau_n f \rightarrow f$ as $n \rightarrow \infty$. It is called a strongly Ditkin set if such a sequence $\{\tau_n\}$ can be chosen uniformly for all functions $f \in I_A(E)$.

In a similar way operator Ditkin sets were defined in [ShT1]. By [ST] the elements of $V^\infty(G)$ are the multipliers of $T(G)$, i.e. $w\omega \subset \omega$ if $w \in V^\infty(G)$ and $\omega \in T(G)$. We call a pseudo-closed subset $E \subset G \times G$ an m -Ditkin set if for any $w \in \Phi(E)$ there exists a sequence $\tau_n \in V^\infty(G)$ such that τ_n vanishes on a pseudo-neighbourhood of E ($n = 1, 2, \dots$) and

$$\|\tau_n w - w\|_{T(G)} \rightarrow 0, \text{ as } n \rightarrow \infty.$$

$E \subset G \times G$ is said to be a strong m -Ditkin set if such a sequence $\{\tau_n\}_{n=1}^\infty$ can be chosen uniformly for all $w \in T(G)$.

If G is a compact metrizable abelian group, it is known that $E = \{0_G\}$ is a strong Ditkin set. Moreover, this set satisfies the following three conditions: (1) E is a set of synthesis; (2) there exist open sets Ω_n containing E such that

$$\Omega_{n+1} \subset \Omega_n \quad n = 1, 2, \dots \text{ and } \bigcap_{n=1}^\infty \bar{\Omega}_n = E;$$

(3) there exists a sequence $\{u_n\}$ with $1 - u_n \in J_A^0(E)$, $n = 1, 2, \dots$, satisfying the following two conditions:

$$\begin{aligned} u_n(x) &= 0 \text{ for all } x \notin \Omega_n, \\ \|u_n\| &\leq 1 + \varepsilon_n, \end{aligned}$$

where $\{\varepsilon_n\}$ is a sequence decreasing to zero.

Note that any closed subset E of the spectrum of a semisimple, regular, commutative Banach algebra A satisfying the conditions (1), (2) and (3) is a strong Ditkin set for A ([D]).

Let Z be a standard Borel space and let $f : G \rightarrow Z$ and $g : G \rightarrow Z$ be Borel functions. Consider

$$(5.11) \quad E = \{(s, t) : f(s) = g(t)\} \subset G \times G.$$

If G, Z are compact metrizable spaces, f, g are continuous functions and $A = V(G) = C(G) \hat{\otimes} C(G)$, the Varopoulos algebra, then E is shown in [D] to satisfy the conditions (1), (2) and (3) and therefore to be a strong Ditkin set for $V(G)$. Knowing that E is a set

of synthesis with respect to m ([ShT1, Theorem 4.8]) we can use a similar argument to show the following statement.

Proposition 5.1. $E = \{(s, t) : f(s) = g(t)\} \subset G \times G$ is a strong m -Ditkin set.

Proof. First we embed Z into the torus, \mathbb{T}_∞ , of infinite dimension: by [T, Theorem A.1] there exists a Borel injective mapping $\psi : Z \rightarrow \mathbb{T}_\infty$. Consider a mapping $\rho : G \times G \rightarrow \mathbb{T}_\infty$ given by

$$\rho(s, t) = \psi(f(t)) - \psi(g(s)),$$

where the subtraction is taken with respect to the group structure on \mathbb{T}_∞ . Then $\rho^{-1}(\mathbb{T}_\infty) = E$.

As $\{0_{\mathbb{T}_\infty}\}$ satisfies (1), (2), (3) in $A(\mathbb{T}_\infty)$, there exist open sets $\Sigma_n \subset \mathbb{T}_\infty$ such that $0_{\mathbb{T}_\infty} \in \Sigma_n$, $\Sigma_n \subset \Sigma_{n+1}$, $n = 1, 2, \dots$, $\bigcap \Sigma_n = \{0\}$ and a sequence of functions $w_n \in A(\mathbb{T}_\infty)$ such that $1 - w_n \in J_A^0(\{0_{\mathbb{T}_\infty}\})$, $w_n(x) = 0$ for $x \notin \Sigma_n$ and $\|w_n\|_A \leq 1 + \varepsilon_n$ ($\varepsilon_n \rightarrow 0$, $\varepsilon_n > 0$).

Define $\Omega_n = \rho^{-1}(\Sigma_n)$. As $\psi \circ f$ and $\psi \circ g$ are Borel mappings for given $m > 0$ there exist by Lusin's theorem closed subsets $A_m \subset G$ and $B_m \subset G$ such that ρ is continuous on $A_m \times B_m$ and $|G \setminus A_m| < 1/m$, $|G \setminus B_m| < 1/m$. We can choose those subsets increasing in m . Therefore

$$\Omega_n \cap (A_m \times B_m) = \{(x, y) \in A_m \times B_m : \rho(x, y) \in \Sigma_n\}$$

is open in $A_m \times B_m$ for all n and

$$\begin{aligned} (5.12) \quad E \cap (A_m \times B_m) &= \bigcap_{n=1}^{\infty} \rho^{-1}(\Sigma_n) \cap (A_m \times B_m) \\ &\subset \bigcap_{n=1}^{\infty} \overline{\Omega_n \cap (A_m \times B_m)} \subset \\ &\subset \bigcap_{n=1}^{\infty} \rho^{-1}(\overline{\Sigma_n}) \cap (A_m \times B_m) = E \cap (A_m \times B_m) \end{aligned}$$

Set $u_n = w_n \circ \rho$. First we show that $u_n \in V^\infty(G)$. In fact, if $w_n = \sum_{\chi \in \hat{\mathbb{T}}_\infty} a_{\chi, n} \chi$ with $\sum_{\chi \in \hat{\mathbb{T}}_\infty} |a_{\chi, n}| = \|w_n\|_A$, then

$$w_n(\rho(s, t)) = \sum_{\chi \in \hat{\mathbb{T}}_\infty} a_{\chi, n} \chi(\psi(f(s))) \overline{\chi(\psi(g(t)))}$$

and our claim follows since

$$\sum_{\chi \in \hat{\mathbb{T}}_\infty} |a_{\chi, n}| |\chi(\psi(f(s)))|^2 = \sum_{\chi \in \hat{\mathbb{T}}_\infty} |a_{\chi, n}| |\chi(\psi(g(t)))|^2 = \sum_{\chi \in \hat{\mathbb{T}}_\infty} |a_{\chi, n}| = \|w_n\|_A.$$

Moreover, by Theorem 3.1, $\|u_n\|_{V^\infty} \leq \|w_n\|_A \leq 1 + \varepsilon_n$, $u_n = 0$ on Ω_n^c and $\tau_n = 1 - u_n$ vanishes on a pseudo-neighbourhood of E .

We next show that for given $w \in \Phi(E)$, $\|\tau_n w - w\| \rightarrow 0$, as $n \rightarrow \infty$. Assume first that $\text{supp}(w) \subset K \times K$, where K is a compact set. Then $w = w_1^m + w_2^m + w_3^m + w_4^m$, where $w_1^m = w \chi_{A_m \times B_m}$, $w_2^m = w \chi_{(G \setminus A_m) \times B_m}$, $w_3^m = w \chi_{A_m \times (G \setminus B_m)}$, $w_4^m = w \chi_{(G \setminus A_m) \times (G \setminus B_m)}$. For given $\varepsilon > 0$ there exists $M > 0$ such that $\|w_i^m\|_{T(G)} < \varepsilon$ for each $m \geq M$ and $i = 2, 3, 4$. Indeed, since the sequence $\{A_m\}$ is increasing in measure to G , by Lebesgue's theorem $\int_{G \setminus A_m} \sum_{i=1}^{\infty} |f_i(r)|^2 dr \rightarrow 0$ as $m \rightarrow \infty$ and for w_2^m we have

$$\|w_2^m\|_{T(G)}^2 \leq \int_{G \setminus A_m} \sum_{i=1}^{\infty} |f_i(r)|^2 dr \int_G \sum_{i=1}^{\infty} |g_i(r)|^2 dr \rightarrow 0.$$

Similarly, $\|w_i^m\|_{T(G)}^2 \rightarrow 0$, $i = 3, 4$. Fix now $m > M$. By [ShT1, Theorem 4.8] E is a set of operator synthesis with respect to Haar measure and by [ShT1, Lemma 6.1] so is $E \cap (A_m \times B_m)$. As $E \cap (A_m \times B_m)$ is closed, by [A, 2.29] there exists $\psi \in T(G)$, $\text{supp } \psi \subset A_m \times B_m$, vanishes on an open neighborhood $\Omega \subset A_m \times B_m$ of $E \cap (A_m \times B_m)$ and $\|w_1^m - \psi\| < \varepsilon$. By (5.12),

$$\bigcap_{n=1}^{\infty} \overline{\Omega_n \cap (A_m \times B_m)} \cap K \times K \subset \Omega \cap K \times K$$

and so $\Omega_n \cap (A_m \times B_m) \cap K \times K \subset \Omega \cap K \times K$ for $n > N$, N is large enough. Thus $u_n \psi \chi_{K \times K} = 0$ for $n > N$. We get

$$\|\tau_n w_1^m - w_1^m\| = \|\tau_n(w_1^m - \psi) \chi_{K \times K} - u_n \psi \chi_{K \times K} - (w_1^m - \psi) \chi_{K \times K}\| \leq (1 + \|\tau_n\|_{V^\infty}) \varepsilon$$

and

$$\|\tau_n w - w\| \leq \|\tau_n w_1^m - w_1^m\| + \sum_{i=2}^4 \|(\tau_n - 1)w_i^m\| \leq 4(1 + \|\tau_n\|_{V^\infty}) \varepsilon.$$

As G is σ -compact there exist compact subsets K_l such that $K_l \subset K_{l+1}$ and $\bigcup_{l=1}^{\infty} K_l = G$, and therefore for any $w \in T(G)$, $w = \lim w \chi_{K_l \times K_l}$ in $T(G)$ giving that $\|\tau_n w - w\| \rightarrow 0$ for any $w \in \Phi(E)$. \square

Corollary 5.2. *Any finite union of sets of type (5.11) is a Ditkin set.*

Proof. It follows from Proposition 5.1 and [ShT1, Theorem 7.1]. \square

Remark 5.3. It follows, in particular, from Corollary 5.2 that finite unions of sets of type (5.11) are sets of operator synthesis. This was also proved by Todorov in [To] using another method.

We will now establish a connection between (strong) Ditkin sets for $A(G)$ and (strong) operator Ditkin sets.

For $w \in T(G)$, define as in [V]

$$Qw(s) = \int_G w(sr, r) dr.$$

If $w = \sum_{i=1}^{\infty} f_i \otimes g_i$ with $\sum_{i=1}^{\infty} \|f_i\|_2^2 \cdot \sum_{i=1}^{\infty} \|g_i\|_2^2 < \infty$. Then

$$Qw(s) = \sum_{i=1}^{\infty} (g_i * \check{f}_i)(s^{-1}) \in A(G)$$

and, moreover, $\|Qw\| \leq \|w\|_{T(G)}$. Thus $Q : T(G) \rightarrow A(G)$ defines a contraction operator.

Theorem 5.4. *Let G be a second countable locally compact group. If E^* is a strong m -Ditkin set then E is a local Ditkin set. If E is a strong Ditkin set then E^* is an m -Ditkin set.*

Proof. Assume first that E^* is a strong m -Ditkin set. Let $\{\Psi_n\}_{n=1}^{\infty} \subset V^\infty(G)$ be a sequence from the definition of a strong m -Ditkin set and let $u \in I_c(E)$. For a compact subset $K \subset G$ containing the support of u , define $(Nu)_K(s, t) = u(st^{-1}) \chi_K(t)$. We have $(Nu)_K(s, t) = \chi_{MK}(s) u(st^{-1}) \chi_K(t)$, where $M = \text{supp}(u)$. As $u(st^{-1}) \in V^\infty(G)$ and $|MK| < \infty$, $|K| < \infty$

∞ , it yields $(Nu)_K \in T(G)$. Moreover, $(Nu)_K$ vanishes on E^* . Therefore, $\|\Psi_n(Nu)_K - (Nu)_K\|_{T(G)} \rightarrow 0$ as $n \rightarrow \infty$. Thus given $\varepsilon > 0$, there exists N such that for $n > N$

$$\|Q(\Psi_n(Nu)_K) - Q((Nu)_K)\|_A \leq \|\Psi_n(Nu)_K - (Nu)_K\|_{T(G)} < \frac{\varepsilon}{|K|}.$$

On the other hand,

$$\begin{aligned} Q(\Psi_n(Nu)_K)(s) &= \int_G \Psi_n(sr, r) Nu(sr, r) \chi_K(r) dr = \int_G \Psi_n(sr, r) u(s) \chi_K(r) dr \\ &= u(s) \int_G \chi_{MK}(sr) \Psi_n(sr, r) \chi_K(r) dr = u(s) Q(\Psi_n(\chi_{MK} \otimes \chi_K))(s) \end{aligned}$$

and similarly $Q((Nu)_K) = u|K|$ giving us

$$\|Q(\Psi_n(Nu)_K) - Q((Nu)_K)\|_A = \|uw_n - u|K|\|,$$

where $w_n = Q(\Psi_n(\chi_M \otimes \chi_K))$. We obtain now that for $\tau_n = w_n/|K|$,

$$\|u\tau_n - u\|_A < \frac{|K|\varepsilon}{|K|} = \varepsilon.$$

Hence E is a local Ditkin set.

Suppose E is a strong Ditkin set. Let $v_n \in J_A^0(G)$ be a sequence such that $\|v_n f - f\|_A \rightarrow 0$ as $n \rightarrow \infty$ for any $f \in I(E)$. By Proposition 3.1, $Nv_n \in V^\infty(G)$, $n = 1, 2, \dots$, and

$$(5.13) \quad \|(Nv_n)(Nf) - Nf\|_{V^\infty} \leq \|v_n f - f\|_A \rightarrow 0.$$

Next we show that $(Nv_n)w \cdot T \rightarrow w \cdot T$ ultra-weakly for any compactly supported T in $B(L_2(G))$ and any compactly supported w in $V^\infty(G)$ such that $w = 0$ on E^* using arguments similar to the ones in the proof of Theorem 4.10.

Assume $\text{supp}(w) \subset K \times K$, for a compact set $K \subset G$. Let w_{kj}^π and \tilde{w}_{kj}^π be V^∞ -functions as defined in (4.5) and let u_{kj}^π be the matrix coefficient $(\pi(\cdot)e_k, e_j)$, $\pi \in \hat{G}$. We have $w_{kj}^\pi(s, t) = \sum_l u_{lj}^\pi(s) \tilde{w}_{kl}^\pi(s, t)$ by (4.7). Hence, for $u \in A(G)$, $\Psi \in T(G)$,

$$\begin{aligned} |\langle (Nv_n - 1)w_{kj}^\pi \cdot T, N(u)\Psi \rangle| &\leq \sum_l |\langle (Nv_n - 1)\tilde{w}_{kl}^\pi \cdot T, u_{jl}^\pi N(u)\Psi \rangle| \\ &= \sum_{l \in L} |\langle (Nv_n - 1)\tilde{w}_{kl}^\pi \cdot T, u_{jl}^\pi N(u)\Psi \rangle| + \sum_{l \notin L} |\langle (Nv_n - 1)\tilde{w}_{kl}^\pi \cdot T, u_{jl}^\pi N(u)\Psi \rangle| \end{aligned}$$

where L is a finite set. For the infinite sum we can apply the estimate (4.9) with $(Nv_n - 1)\Psi$ instead of Ψ , so that for given $\varepsilon > 0$ there exists a finite subset L such that

$$\sum_{l \notin L} |\langle (Nv_n - 1)\tilde{w}_{kl}^\pi \cdot T, u_{jl}^\pi N(u)\Psi \rangle| = \sum_{l \notin L} |\langle \tilde{w}_{kl}^\pi \cdot T, u_{jl}^\pi N(u)(Nv_n - 1)\Psi \rangle| < \varepsilon$$

For the finite sum we note first that $\tilde{w}_{kl}^\pi = N(v_{kl}^\pi)$ for some $v_{kl}^\pi \in M_{cb}A(G)$ by (4.10). Then by (5.13) there exists $M > 0$ such that for any $n > M$,

$$\begin{aligned} \sum_{l \in L} |\langle (Nv_n - 1)\tilde{w}_{kl}^\pi \cdot T, u_{jl}^\pi N(u)\Psi \rangle| &= \sum_{l \in L} |\langle T, u_{jl}^\pi (Nv_n - 1)N(v_{kl}^\pi)N(u)\Psi \rangle| = \\ \sum_{l \in L} |\langle T, u_{jl}^\pi (Nv_n - 1)N(v_{kl}^\pi u)\Psi \rangle| &\leq \sum_{l \in L} \|T\| \cdot \|(Nv_n - 1)N(v_{kl}^\pi u)\|_{V^\infty} \|\Psi\|_{T(G)} < \varepsilon \end{aligned}$$

We obtain,

$$\langle (Nv_n - 1)w_{kj}^\pi \cdot T, N(u)\Psi \rangle \rightarrow 0, \quad n \rightarrow \infty.$$

As in the proof of Theorem 4.10, it yields

$$(5.14) \quad \langle (Nv_n - 1)w_{kj}^\pi \cdot T, \Psi \rangle \rightarrow 0, \quad n \rightarrow \infty$$

for any $\Psi \in T(G)$, since $\{Nv_n\}$ is bounded in norm. Using the same arguments as in the end of the proof of Theorem 4.3 we have that for big enough compact set K there exists a linear combination $u = \sum_{i=1}^d c_i u_i$ of matrix coefficients such that

$$\|w - u\chi_{K^{-1}K} \cdot w\|_{T(G)} < \varepsilon.$$

Thus for $\omega \in T(G)$, $\text{supp}(\omega) \subset K \times K$,

$$\begin{aligned} |\langle (Nv_n - 1)w \cdot T, \omega \rangle| &\leq |\langle (Nv_n - 1) \cdot T, \omega(w - u\chi_{K^{-1}K} \cdot w) \rangle| \\ &\quad + |\langle (Nv_n - 1) \cdot T, \omega(u\chi_{K^{-1}K} \cdot w) \rangle|. \end{aligned}$$

The first summand is less than $C\varepsilon$ for some constant C , for n large enough, as $\{Nv_n\}$ is bounded in norm. As $\omega(u_{kj}^\pi \cdot w) = \omega w_{kj}^\pi$ by (4.5) for all k, j, π , we have by (5.14)

$$\langle (Nv_n - 1) \cdot T, \omega(u_{kj}^\pi \chi_{K^{-1}K} \cdot w) \rangle = \langle (Nv_n - 1)w_{kj}^\pi \cdot T, \omega \rangle \rightarrow 0, \quad n \rightarrow \infty,$$

and hence $|\langle (Nv_n - 1) \cdot T, \omega(u\chi_{K^{-1}K} \cdot w) \rangle| < \varepsilon$ for n large enough. Thus,

$$\langle (Nv_n - 1)w \cdot T, \omega \rangle \rightarrow 0$$

for each $\omega \in T(G)$. Hence $(Nv_n)w\omega \rightarrow w\omega$ weakly for each $\omega \in T(G)$. As the set of all linear combinations of $w\omega$, where $w \in V^\infty$, $w = 0$ on E^* and $\omega \in T(G)$, is dense in $\Phi(E)$ ([**ShT2**, Proposition 5.3]), $(Nv_n)\omega \rightarrow \omega$ weakly for any $\omega \in \Phi(E)$. Taking if necessary convex linear combinations of the Nv_n 's, we obtain elements $w_n \in V^\infty(G)$, $n \in \mathbb{N}$, such that $\|w_n\omega - \omega\|_{T(G)} \rightarrow 0$ as n tends to infinity. Clearly, the sequence $\{w_n\}_n$ satisfies the necessary conditions. \square

Corollary 5.5. *Any closed subgroup H of G is a local Ditkin set.*

Proof. We have $H^* = \{(s, t) : f(s) = f(t)\}$, where $f : G \rightarrow H \backslash G$, $t \mapsto Ht$. As G is metrizable, by [**HRI**, ((8.14))], $H \backslash G$ is metrizable. The statement now follows from Proposition 5.1 and Theorem 5.4. \square

For G amenable the statement was obtained in [**FKLS**] showing that $I_A(H)$ has a bounded approximate identity. For arbitrary locally compact G and H a closed neutral subgroup it was proved in [**DD**]. Note that there are closed subgroups of l.c.s.c. groups which are not neutral (see [**RD**, p.107]).

Acknowledgments.

We are indebted to A.T.Lau, V.Shulman and N.Spronk for helpful discussions and valuable information.

References

- [A] W.Arveson, Operator algebras and invariant subspaces. *Ann. of Math. (2)* **100** (1974), 433–532.
- [BSm] D.P.Blecher, R.R.Smith, The dual of the Haagerup tensor product. *J. London Math. Soc. (2)* **45** (1992), no.1, 126–144.
- [Ca] J. de Canniere, Unbounded approximate units of the Fourier algebra of $SO_e(n, 1)$. *Operator algebras and group representations*, Proc. int. Conf., Neptun/Rom. 1980, Vol. I, Monogr. Stud. Math. 17, 143-146 (1984).
- [CaH] J. de Canniere and U.Haagerup, Multipliers of the Fourier algebras of some simple Lie groups and their discrete subgroups. *Amer. J. Math.*, **107** (1984), 455-500.
- [CoH] M.Cowling and U.Haagerup, Completely bounded multipliers of the Fourier algebra of a simple Lie group of real rank 1. *Invent. Math.*, **96** (1989), 507-549.
- [DD] J.Delaporte, A.Derighetti, Invariant projections and convolution operators. *Proc. Amer. Math. Soc.* **129** (2001), no. 5, 1427–1435.
- [Di] J. Dixmier, *Les C^* -algèbres et leurs représentations*. 2e ed. Cahiers scientifiques. 29. Paris: Gauthier-Villars. XV, 390 p. (1969).
- [D] S.W. Drury, On non-triangular sets in tensor algebras. *Studia Math.* **34** 1970 253–263.
- [EfR] E.G.Effros and Z.J.Ruan, *Operator spaces*. London Mathematical Society Monographs. New Series, 23. The Clarendon Press, Oxford University Press, New York, 2000. xvi+363 pp.
- [ErKSh] J.A.Erdos, A.Katavolos, V.S.Shulman, Rank one subspaces of bimodules over maximal abelian selfadjoint algebras. *J. Funct. Anal.* **157** (1998), no. 2, 554–587.
- [E] P.Eymard, L’algèbre de Fourier d’un groupe localement compact. (French) *Bull. Soc. Math. France* **92** 1964 181–236.
- [FKLS] B.Forrest, E.Kaniuth, A.T.Lau, N.Spronk, Ideals with bounded approximate identities in Fourier algebras. *J. Funct. Anal.* **203** (2003), no. 1, 286–304.
- [F] J.Froelich, Compact operators, invariant subspaces, and spectral synthesis. *J. Funct. Anal.* **81** (1988), no. 1, 1–37.
- [He] C.Herz, Harmonic synthesis for subgroups. *Ann. Inst. Fourier (Grenoble)* **23** (1973), no. 3, 91–123.
- [HRI] E.Hewitt, K.A.Ross, *Abstract harmonic analysis*. Vol. I. Second edition. Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], 115. Springer-Verlag, Berlin-New York, 1979. ix+519.
- [HRII] E.Hewitt, K.A.Ross, *Abstract harmonic analysis*. Vol. II. Die Grundlehren der mathematischen Wissenschaften, Band 152 Springer-Verlag, New York-Berlin 1970 ix+771 pp.
- [RD] W.Roelcke, S.Dierolf, *Uniform Structures on Topological Groups and their Quotients*, McGraw-Hill International book Company, 1981, pp. 276.
- [ShT1] V.Shulman, L. Turowska, Operator synthesis. I. Synthetic sets, bilattices and tensor algebras. *J. Funct. Anal.* **209** (2004), no. 2, 293–331.
- [ShT2] V.Shulman and L.Turowska, Operator synthesis II. Individual operator synthesis and linear operator equations, Preprint 2004, Department of Mathematics, Chalmers University of Technology and Göteborg University, Sweden.
- [Sm] R.R. Smith, Completely bounded module maps and the Haagerup tensor product, *J.Funct.Anal.* **102** (1991), 156-175.
- [S] N.Spronk, On multipliers of the Fourier algebra of a locally compact group, PhD Thesis, University of Waterloo.
- [ST] N.Spronk and L.Turowska, Spectral synthesis and operator synthesis for compact groups. *J. London Math. Soc. (2)* **66** (2002), no. 2, 361–376.
- [T] M.Takesaki, *Theory of operator algebras I*. Springer-Verlag, 1979.
- [TT] M.Takesaki, N.Tatsuuma, Duality and subgroups. II. *J. Functional Analysis* **11** (1972), 184–190.

- [To] I.Todorov, Synthetic properties of ternary masa-bimodules, to appear in Houston J. Math.
[V] N.Th.Varopoulos, Tensor algebras and harmonic analysis, Acta Math. **119** (1967), 51–112.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF METZ,
F-57045 METZ, FRANCE
E-mail address: ludwig@poncelet.sciences.univ-metz.fr

DEPARTMENT OF MATHEMATICS, CHALMERS UNIVERSITY OF TECHNOLOGY,
SE-412 96 GÖTEBORG, SWEDEN
E-mail address: turowska@math.chalmers.se

The work was partially written when the first author was visiting Chalmers University of Technology in Göteborg, Sweden and when the second author was a visiting professor at Metz University, France. The first author was supported by the research grants CUL/01/014 and R1F104C09 of the University of Luxembourg and the second author was also supported by the Swedish Research Council.